



Instructions for Use

The following Instructions for Use should be followed when starting and continuing on your treatment:

Meet your KESIMPTA® Sensoready® Pen



No visible needle

- Press against skin to start

Green progress indicator

- Stops moving when it's complete

Can be kept at room temperature for up to a week

- It's travel friendly!

2 clicks

- 1 click when it starts
- 1 click when it's almost done

Your dosing schedule²

Each pre-filled Sensoready® pen contains 20 mg KESIMPTA®.

- Take your initial dose (1st dose) under the guidance of a healthcare professional
- Take the 2nd dose on the same day of the second week (week 1)
- Take the 3rd dose on the same day of the third week (week 2)
- After the first 3 injections, do not inject the following week (week 3)
- Start your monthly doses on week 4, taking your medication on the same date every month moving forward.



Once you start, treatment can be straightforward to fit into your routine.

Tracking Tip:
Keep it simple, take your medication on the same date every month.



Using your KESIMPTA® Sensoready® pen



STEP 1: Preparation

**Take the pen out of the fridge
15 to 30 minutes before
injection to allow it to reach
room temperature.**

See page
18 for help
picking an
injection
site

- Look through the viewing window. The liquid should be clear to slightly opalescent and colourless to slightly brownish-yellow. You may see a small air bubble, which is normal.
- Check the expiry date of the pen (found on the outer carton or the label on the pen).




Important information:

- Do not use if the liquid has visible particles or is cloudy.
- Do not use if the pen has expired.
- **Contact your pharmacist or healthcare professional if your pen fails any of these checks**

STEP 2: Get ready


Pick and clean your injection site.

- Wash your hands with soap and water.
- Using a circular motion, clean the injection site with the alcohol wipe.
- Let it dry completely.
- Do not touch the injection site again before injecting



Remove the cap and throw it away.


- Twist off the cap in the direction of the arrow. Don't try to reattach it.
- Make sure you use the pen within 5 minutes of removing the cap and do not shake.
- You may see a few drops of medicine come out of the needle. This is normal.



STEP 3: Inject


Hold the pen at a 90° angle to the cleaned injection site.

- **Press down** to activate the pen and **hold**. You'll hear the first click and the green indicator will start moving. Keep the pen pressed firmly against your skin.



Watch the green indicator fill.

- Listen for the second click to signal you're almost done. Once the green indicator is full and has stopped, you'll know that you've done it right and you can now remove the pen.
- If there's blood at the injection site, press down with a cotton ball/gauze, but don't rub.



Important information:

If the green indicator does not fill the window, this means you have not received the full dose. Contact your doctor or pharmacist if the green indicator is not visible.

STEP 4: Finish

Dispose of the used pen in a sharps bin

(i.e. a puncture-resistant closable container or similar), if provided.

Please ask your doctor or pharmacist for more information on how to dispose of your used pens safely.




Important information:

- Never try to re-use your pen.
- Always keep the sharps container out of reach of children
- Do not throw away any medicines via wastewater or household waste (including flushing down the toilet or putting down the sink).

Choosing the appropriate spot

The front of the thigh is the recommended site.²

Change your injection spot each month



Outer upper arm

The outer upper arm, may be used by a healthcare professional or caregiver if they are injecting you.²

Lower abdomen (stomach area)

The lower abdomen (stomach area) except the area 5 cm around your belly button.²

Thigh

The front of the thigh is the recommended site.²



X DO NOT inject into areas where the skin is tender, bruised, red, scaly or hard. Avoid areas with scars, stretch marks or infection sites.²

Important safety checks



Do not use this medicine:²

- If either the seal on the outer package or the seal on the pen is broken. Keep the pen in the sealed outer package until you are ready to use it.
- If you drop your pen, do not use it if the pen looks damaged, or if you dropped it with the cap removed.
- After the expiry date, which is stated on the outer box or the label on the pen after 'EXP'. The expiry date refers to the last day of that month.
- If the solution contains visible particles or is cloudy.



Tracking tip

You should make a note of when and where on your body you inject. There's a handy tracker at the back of this book for you to use to do this.

Forgot a dose in your first month?

Administer your dose as soon as you remember and time your future injections from the day you injected this dose.²

